

USAGE AREA	APPLICATION TIME	SOIL APP.	FOLIAR APP.
GREENHOUSE AND OUTDOOR VEGETABLES (Tomato, Pepper, Cucumber, Eggplant, Bean, Strawberry etc.)	at intervals of 15-20 days after the first fruits	300 gr / Da	100 lt water 150-200 Gr
WINTER VEGETABLES WITH LEAVES (Curly, Lettuce, Cabbage, Spinach, etc.)	With an interval of 15-20 days from the period when it has 3-4 leaves	300 gr / Da	100 lt water 150-200 Gr
BULK PLANTS (Melon, Watermelon, Pumpkin, Potato, Onion etc.)	15-20 days from the 4-6 leaf stage and after the first fruits	300 gr / Da	100 lt water 150-200 Gr
ALL FRUIT TREES (Peach, Apricot, Cherry, Plum Apple, Pear, Quince, Citrus, Olive, Vineyard, Hazelnut, Pistachio etc.)	It is applied in two or three repetitions with the start of development.	0,5-1 Kg/Da	100 lt water 150-200 Gr
INDUSTRIAL PLANTS (Corn, Cotton, Sunflower etc.)	With an interval of 15-20 days starting from the period with 5-6 leaves	300 gr / Da	100 lt water 150-200 Gr
ALL LEGUMES (Chickpea, Bean Lentil, Soybean Sunflower etc.)	5-6 leaf period and 15-20 days later	300 gr / Da	100 lt water 150-200 Gr
GRAINS (Wheat, Barley, Paddy etc.)	During tillering period	300 gr / Da	100 lt water 150-200 Gr
Green Areas	Throughout the development period	300 gr / Da	100 lt water 150-200 Gr