

USAGE AREA	APPLICATION TIME	SOIL APPLICATION	FOLIAR APPLICATION
All Greenhouse and All Open Field Vegetables	1 week after germination. First flowering. After the first fruit formation.	500 - 1 kg / Ha	100- 150 Gr./ 100 Lt. Water
All Stone and Pome Fruit Trees	At the beginning of flowering. 15 days after fruit formation. 30 days before the harvest.	500 - 1 kg / Ha	100- 150 Gr./ 100 Lt. Water
Citrus, Banana, Fig and Olive	At the beginning of flowering. 15 days after fruit formation. 30 days before the harvest.	500 - 1 kg / Ha	100-150 Gr./ 100 Lt. Water
Vineyards and Ornamental Plants	At the beginning of flowering. 15 days after fruit formation. 30 days before the harvest.	500 -1 kg / Ha	100- 150 Gr./ 100 Lt. Water
Corn and Sunflower	First Application: 30 days after sowing. Second Application: as required.	500 -1 kg / Ha	100- 150 Gr./ 100 Lt. Water
Cotton	From the beginning of first square - bud, it can be applied at regular intervals when needed.	500 - 1 kg / Ha	100- 150 Gr./ 100 Lt. Water
Industrial Plants (Soya, Peanut and Canola)	After the 3-5 leaf growth stage, it can be applied at regular intervals when needed.	500 - 1 kg / Ha	100- 150 Gr./ 100 Lt. Water
Field crops (Wheat, Barley, Rice) and Green Areas (Golf Courses, Lawn Care and Sports Turf)	When elongation of field crops reaches 20 - 25 cm. The first period of kernel development (watery ripe and milk stages).	500 - 1 kg / Ha	100- 150 Gr. / 100 Lt. Water