

USAGE AREA	APPLICATION TIME
<b>All greenhouse and all open field and winter vegetables</b>	1-2 app.with drip irrigation 10 days after planting. From pre-flowering until the harvest, it can be applied with foliar app.at regular intervals
<b>Melon, watermelon, zucchini, cauliflower, potato, onion, radish, carrot and garlic</b>	From planting until the beginning of ripening 2-3 app.
<b>Strawberry</b>	From the vegetative period until the harvesting 3-4 app.
<b>Ornamental plants</b>	Begins during vegetative growth and apply for 10-15 day intervals throughout the season.
<b>Vineyard</b>	From the beginning of the shoot until the harvesting period, it can be applied by drip irrigation when needed. It can also be applied by foliar app.at regular intervals until fruit set period.
<b>Corn and sunflower</b>	After planting, it is recommended that the product is applied after 15 days.
<b>Banana</b>	2-3 applications from 30-45 days prior to blooming(april-may)
<b>Cereals and forage crops</b>	During the tillering period in early spring.
<b>Apple, pear, cherry, sour cherry, apricot, peach, almond, hazelnut,</b>	From beginning of blooming until harvesting, it can be applied at regular intervals.
<b>Olive, fig and citrus</b>	From pre-flowering until the harvest period, 2-3 applications.

<b>SOIL APP.</b>	<b>FOLIAR APP. (100 lt water)</b>
10-20 lt/ha	150-200 cc
10-20 lt/ha	150-200 cc
10-20 lt/ha	150-200 cc
10-20 lt/ha	150-200 cc
10-20 lt/ha	150-200 cc
10-20 lt/ha	200-250 cc
10-20 lt/ha	200-250 cc
10-20 lt/ha	250-300 cc
10-20 lt/ha	200-250 cc
5-10 lt/ha	200-250 cc